

others; by Kunze.—On *Conyza chilensis*, Spreng, and *C. diversifolia*, Weinw.; by Weinmann.—Review of a century of Brazilian *Malpighiaceæ*; by Dr. Grisebach.—On Mexican Plants collected by Schiede, &c.; by Schlechtendal.

Plantes Cryptogames de France. Par J. B. H. Q. Desmazières.

Of this valuable work three Numbers, 18—20, have appeared since our last notice. It would take up too much room to give a list of the species, which amount to 150, contained in them; but it may truly be said that with every number the interest increases.

Deutschlands Kryptogamische Giftgewächse, &c. Von Dr. P. Phœbus. Berlin, 1838. 4to.

This is a continuation of Brandt and Ratzeburg's account of the poisonous plants of Germany. With the exception of the Fungi, the cryptogamic plants noticed as possessing deleterious qualities are confined to one or two species of *Equisetum* and *Lycopodium*. The poisonous fungi are admirably described, with remarkably full synonyms, and of many of the more common ones there are excellent figures, with a very correct analysis. There is a very complete account of Ergot beautifully illustrated. We recommend the work very highly to the notice of such as wish to become acquainted with dangerous Fungi, and to the general cryptogamic student. The price is very reasonable.

Fauna Coleopterorum Helvetica, auctore Oswaldo Heer. Pars I. Fasc. I. Turici.—The present first fasciculus contains the *Carabidæ* and the commencement of the *Dytiscidæ*.

We are glad to learn that the Rev. Dr. Pye Smith's Congregational Lecture on the Relation between the Holy Scriptures and some parts of Geological Science is in the press.

PROCEEDINGS OF LEARNED SOCIETIES.

BRITISH ASSOCIATION.

August 26.—J. E. Gray, Esq., V.P., in the Chair.

Secretaries:—Mr. E. Forbes and Mr. Patterson.

The first paper read was by Mr. Lankester, on the Formation of Woody Tissue, the conclusions in which were opposed to the views, supported in this country by Dr. Lindley, of the formation of fibrous tissue by descent from the buds and leaves, and contained some in-

others; by Kunze.—On *Conyza chilensis*, Spreng, and *C. diversifolia*, Weinw.; by Weinmann.—Review of a century of Brazilian *Malpighiaceæ*; by Dr. Grisebach.—On Mexican Plants collected by Schiede, &c.; by Schlechtendal.

Plantes Cryptogames de France. Par J. B. H. Q. Desmazières.

Of this valuable work three Numbers, 18—20, have appeared since our last notice. It would take up too much room to give a list of the species, which amount to 150, contained in them; but it may truly be said that with every number the interest increases.

Deutschlands Kryptogamische Giftgewächse, &c. Von Dr. P. Phœbus. Berlin, 1838. 4to.

This is a continuation of Brandt and Ratzeburg's account of the poisonous plants of Germany. With the exception of the Fungi, the cryptogamic plants noticed as possessing deleterious qualities are confined to one or two species of *Equisetum* and *Lycopodium*. The poisonous fungi are admirably described, with remarkably full synonyms, and of many of the more common ones there are excellent figures, with a very correct analysis. There is a very complete account of Ergot beautifully illustrated. We recommend the work very highly to the notice of such as wish to become acquainted with dangerous Fungi, and to the general cryptogamic student. The price is very reasonable.

Fauna Coleopterorum Helvetica, auctore Oswaldo Heer. Pars I. Fasc. I. Turici.—The present first fasciculus contains the *Carabidæ* and the commencement of the *Dytiscidæ*.

We are glad to learn that the Rev. Dr. Pye Smith's Congregational Lecture on the Relation between the Holy Scriptures and some parts of Geological Science is in the press.

PROCEEDINGS OF LEARNED SOCIETIES.

BRITISH ASSOCIATION.

August 26.—J. E. Gray, Esq., V.P., in the Chair.

Secretaries:—Mr. E. Forbes and Mr. Patterson.

The first paper read was by Mr. Lankester, on the Formation of Woody Tissue, the conclusions in which were opposed to the views, supported in this country by Dr. Lindley, of the formation of fibrous tissue by descent from the buds and leaves, and contained some in-

others; by Kunze.—On *Conyza chilensis*, Spreng, and *C. diversifolia*, Weinw.; by Weinmann.—Review of a century of Brazilian *Malpighiaceæ*; by Dr. Grisebach.—On Mexican Plants collected by Schiede, &c.; by Schlechtendal.

Plantes Cryptogames de France. Par J. B. H. Q. Desmazières.

Of this valuable work three Numbers, 18—20, have appeared since our last notice. It would take up too much room to give a list of the species, which amount to 150, contained in them; but it may truly be said that with every number the interest increases.

Deutschlands Kryptogamische Giftgewächse, &c. Von Dr. P. Phœbus. Berlin, 1838. 4to.

This is a continuation of Brandt and Ratzeburg's account of the poisonous plants of Germany. With the exception of the Fungi, the cryptogamic plants noticed as possessing deleterious qualities are confined to one or two species of *Equisetum* and *Lycopodium*. The poisonous fungi are admirably described, with remarkably full synonyms, and of many of the more common ones there are excellent figures, with a very correct analysis. There is a very complete account of Ergot beautifully illustrated. We recommend the work very highly to the notice of such as wish to become acquainted with dangerous Fungi, and to the general cryptogamic student. The price is very reasonable.

Fauna Coleopterorum Helvetica, auctore Oswaldo Heer. Pars I. Fasc. I. Turici.—The present first fasciculus contains the *Carabidæ* and the commencement of the *Dytiscidæ*.

We are glad to learn that the Rev. Dr. Pye Smith's Congregational Lecture on the Relation between the Holy Scriptures and some parts of Geological Science is in the press.

PROCEEDINGS OF LEARNED SOCIETIES.

BRITISH ASSOCIATION.

August 26.—J. E. Gray, Esq., V.P., in the Chair.

Secretaries:—Mr. E. Forbes and Mr. Patterson.

The first paper read was by Mr. Lankester, on the Formation of Woody Tissue, the conclusions in which were opposed to the views, supported in this country by Dr. Lindley, of the formation of fibrous tissue by descent from the buds and leaves, and contained some in-

others; by Kunze.—On *Conyza chilensis*, Spreng, and *C. diversifolia*, Weinw.; by Weinmann.—Review of a century of Brazilian *Malpighiaceæ*; by Dr. Grisebach.—On Mexican Plants collected by Schiede, &c.; by Schlechtendal.

Plantes Cryptogames de France. Par J. B. H. Q. Desmazières.

Of this valuable work three Numbers, 18—20, have appeared since our last notice. It would take up too much room to give a list of the species, which amount to 150, contained in them; but it may truly be said that with every number the interest increases.

Deutschlands Kryptogamische Giftgewächse, &c. Von Dr. P. Phœbus. Berlin, 1838. 4to.

This is a continuation of Brandt and Ratzeburg's account of the poisonous plants of Germany. With the exception of the Fungi, the cryptogamic plants noticed as possessing deleterious qualities are confined to one or two species of *Equisetum* and *Lycopodium*. The poisonous fungi are admirably described, with remarkably full synonyms, and of many of the more common ones there are excellent figures, with a very correct analysis. There is a very complete account of Ergot beautifully illustrated. We recommend the work very highly to the notice of such as wish to become acquainted with dangerous Fungi, and to the general cryptogamic student. The price is very reasonable.

Fauna Coleopterorum Helvetica, auctore Oswaldo Heer. Pars I. Fasc. I. Turici.—The present first fasciculus contains the *Carabidæ* and the commencement of the *Dytiscidæ*.

We are glad to learn that the Rev. Dr. Pye Smith's Congregational Lecture on the Relation between the Holy Scriptures and some parts of Geological Science is in the press.

PROCEEDINGS OF LEARNED SOCIETIES.

BRITISH ASSOCIATION.

August 26.—J. E. Gray, Esq., V.P., in the Chair.

Secretaries:—Mr. E. Forbes and Mr. Patterson.

The first paper read was by Mr. Lankester, on the Formation of Woody Tissue, the conclusions in which were opposed to the views, supported in this country by Dr. Lindley, of the formation of fibrous tissue by descent from the buds and leaves, and contained some in-